

LAB 23 – Metabolic Rates

Introduction:


The cardiovascular (circulatory) system functions to deliver oxygen and nutrients to tissues for growth and metabolism, and to remove metabolic wastes. The heart pumps blood through a circuit that includes arteries, arterioles, capillaries, venules, and veins. One important circuit is the pulmonary circuit, where there is an exchange of gases within the alveoli of the lungs. The right side of the human heart receives deoxygenated blood from body tissues and pumps it to the lungs. The left side of the heart receives oxygenated blood from the lungs and pumps it to the tissues.

With increased exercise, several changes occur within the circulatory system, thus increasing the delivery of oxygen to actively respiring muscles cells. These changes include increased heart rate, increased blood flow to muscular tissue, decreased blood flow to nonmuscular tissue, increased arterial pressure, increased body temperature, and increased breathing rate.

Blood Pressure

An important measurable aspect of the circulatory system is blood pressure. When the ventricles of the heart contract, pressure is increased throughout all the arteries. Arterial blood pressure is directly dependent on the amount of blood pumped by the heart per minute and the resistance to blood flow through the arterioles. The arterial blood pressure is determined using a device known as a **sphygmomanometer**. This device consists of an inflatable cuff connected by rubber hoses to a hand pump and to a pressure gauge graduated in millimeters of mercury. The cuff is wrapped around the upper arm and inflated to a pressure that will shut off the brachial artery. The examiner listens for the sounds of blood flow in the brachial artery by placing the bell of a stethoscope in the inside of the elbow below the biceps.

Figure 1



At rest the blood normally goes through the arteries so that the blood in the central part of the artery moves faster than the blood in the peripheral part. Under these conditions, the artery is silent when one listens. When the sphygmomanometer cuff is inflated to a pressure above the systolic pressure, the flow of blood is stopped and the artery is again silent. As the pressure in the cuff gradually drops to levels between the systolic and diastolic pressures of the artery, the blood is pushed through the compressed walls of the artery in a turbulent flow. Under these conditions, the blood is mixed, and the turbulence sets up vibrations in the artery that are

heard as sound in the stethoscope. These sounds are known as the heart sounds, or sounds of Korotkoff. These sounds are divided into five phases based on the loudness and quality of the sounds.

Phase 1. A loud, clear, snapping sound is evident which increases in intensity as the cuff is deflated. In the example shown in **Figure 2** this phase begins at a cuff pressure of 120 millimeters of mercury (mm Hg) and ends at a pressure of 106 mm Hg.

Phase 2. A succession of murmurs can be heard. Sometimes the sounds seem to disappear during this time, which may be a result of inflating or deflating the cuff too slowly. In the example shown in **Figure 2**, this phase begins at a cuff pressure of 106 mm Hg and ends at a pressure of 86 mm Hg.

Phase 3. A loud, thumping sound, similar to that in Phase 1, but less clear, replaces the murmurs. In the example shown in **Figure 2**, Phase 3 begins at a cuff pressure of 86 mm Hg and ends at a pressure of 81 mm Hg.

Phase 4. A muffled sound abruptly replaces the thumping sounds of Phase 3. In the example shown in **Figure 2**, this phase begins at a cuff pressure of 81 mm Hg and ends at a pressure of 76 mm Hg.

Phase 5. All sounds disappear.




Figure 2

The cuff pressure at which the first sound is heard (that is, the beginning of Phase 1) is taken as the systolic pressure. The cuff pressure at which the muffled sound of Phase 4 disappears (the beginning of Phase 5) is taken as the measurement of the diastolic pressure. In the example shown in **Figure 2**, the pressure would be recorded in this example as 120/76. A normal blood pressure measurement for a given individual depends of the person's age, sex, heredity, and environment. When these factors are taken into account, blood pressure measurements that are chronically elevated may indicate a state deleterious to the health of the person. This condition is called hypertension and is a major contributing factor in heart disease and stroke. Typical blood pressure for men and women varies with age and fitness (**Figure 3**). For high school students, the typical range is usually 100-130/70-90.

Figure 3: Typical Blood Pressure for Men and Women at Different Ages

Age (years)	Systolic Pressure		Diastolic Pressure	
	Men	Women	Men	Women
10	103	103	69	70
11	104	104	70	71
12	106	106	71	72
13	108	108	72	73
14	110	110	73	74
15	112	112	75	76
16	118	116	73	72
17	121	116	74	72
18	120	116	74	72
19	122	115	75	71
20-24	123	116	76	72
25-29	125	117	78	74
30-34	126	120	79	75
35-39	127	124	80	78
40-44	129	127	81	80
45-49	130	131	82	82
50-54	135	137	83	84
55-59	138	139	84	84
60-64	142	144	85	85
65-69	143	154	83	85
70-74	145	159	82	85

Activity 1 – Heart Rate and Temperature

In ectothermic animals there is a direct relationship between the rate of many physiological activities and environmental temperature. The rate of metabolism in these animals increases as environmental temperatures increase from approximately 5° C to 35° C. Increasing the temperature by approximately 10° results in a doubling of the metabolic rate. That is why the snake or the lizard can hardly move when it is cold but becomes quite active after warming in the sun.

In this exercise, you will observe a species of Japanese killifish, the Medaka (*Oryzias latipes*) over a range of increasing environmental temperature. You will be able to observe the two-chambered heart beat over the duration of the investigation

Procedure:

1. Watch the video of the Medaka. You should be able to identify the two-chambered heart.
2. To obtain the heart rate in beats per minute, count the heart rate for 15 seconds then multiply by 4.
3. In **Table 1**, record the temperature and heart rate for every change in temperature until the end of the video.

Table 1: Temperature and Heart Rate Data

Temperature	Heart Rate (beats/min)	Temperature	Heart Rate (beats/min)

Activity 2 – Measuring Blood Pressure

Note: These labs are ONLY for experimental, and not diagnostic, purposes.

A sphygmomanometer (blood pressure cuff) is used to measure blood pressure. The cuff, designed to fit around the upper arm, can be expanded by pumping a rubber bulb connected to the cuff. The pressure gauge, scaled in millimeters, indicated the pressure inside the cuff. A stethoscope is used to listen to the individual's pulse (**Figure 1**). The earpieces of the stethoscope should be cleaned with alcohol swabs before and after each use.

Procedure:

1. Work in pairs. Those who are to have their blood pressure measured should be seated with both shirt sleeves rolled up.
2. Attach the cuff of the sphygmomanometer snugly around the upper arm.
3. Place the stethoscope directly below the cuff in the bend of the elbow joint.
4. Close the valve of the bulb by turning it clockwise. Pump air into the cuff until the pressure gauge reaches 180 mm Hg.
5. Turn the valve of the bulb counterclockwise and slowly release air from the cuff. Listen for a pulse.
6. When you first hear the heart sounds, note the pressure on the gauge. This is the systolic pressure.
7. Continue to slowly release air and listen until the clear thumping sound of the pulse becomes strong and then fades. When you last hear the full heart beat, note the pressure. This is the diastolic pressure.
8. Repeat the measurement two more times and determine the average systolic and diastolic pressure, then record these values in the blood pressure data table.
9. Trade places with your partner. When your average systolic and diastolic pressure has been determined, record these values in **Table 2** below.

Table 2: Blood Pressure Data

Measurement	1	2	3	Average
Systolic				
Diastolic				

Activity 3 – A Test of Fitness

The point scores on the following tests provide an evaluation of fitness based not only on cardiac muscular development but also on the ability of the cardiovascular system to respond to sudden changes in demand. **CAUTION: Make sure that you do not attempt this exercise if strenuous activity will aggravate a health problem.** Work in pairs. Determine the fitness level for one member of the pair (Tests 1 → 5) and then repeat the process for the other member. Record all data in the spaces provided and in the Fitness Data Box (**Table 9**) at the end of this exercise **FOR YOUR OWN DATA!**

Test #1: Standing Systolic Compared with Reclining Systolic

Use the sphygmomanometer as you did in Exercise 1 to measure the change in systolic blood pressure from a reclining to a standing position.

Procedure:

1. The subject should recline on a lab table for at least 5 minutes. At the end of this time, measure the systolic and diastolic pressure and record these values below.

reclining systolic pressure

_____ mm Hg

reclining diastolic pressure

_____ mm Hg

2. Remain reclining for two additional minutes, then stand and *immediately* repeat the measurements on the same subject (arms down). Record these values below.

standing systolic pressure

_____ mm Hg

standing diastolic pressure

_____ mm Hg

3. Determine the change in systolic pressure from reclining to standing by subtracting the standing measurement from the reclining measurement. Assign fitness points based on **Table 3** and record in the fitness data box (**Table 9**)

Table 1: Change in Systolic Pressure from Reclining to Standing

Change (mm Hg)	Fitness Points
rise of 8 or more	3
rise of 2-7	2
no rise	1
fall of 2-5	0
fall of 6 or more	-1

Cardiac Rate and Physical Fitness

During physical exertion, the cardiac rate (beats per minute) increases. This increase can be measured as an increase in pulse rate. Although the maximum cardiac rate is generally the same in people of the same age group, those who are physically fit have a higher stroke volume (milliliters per beat) than more sedentary individuals. A person who is in poor physical condition, therefore, reaches his or her maximum cardiac rate at a lower work level than a person of comparable age who is in better shape. Individuals who are in good physical condition can deliver more oxygen to their muscles (have a higher aerobic capacity) before reaching maximum cardiac rate than can those in poor condition.

Thus, the physically fit have a slower increase in their cardiac rate with exercise and a faster return to the resting cardiac rate after exercise. Physical fitness, therefore, involves not only muscular development but also the ability of the cardiovascular system to respond to sudden changes in demand.

Test #2: Standing Pulse Rate

Procedure:

1. The subject should stand at ease for 2 minutes after Test 1.
2. After the 2 minutes, determine the subject's pulse by using the radial artery.
3. Count the number of beats for 30 seconds and multiply by 2. The pulse rate is the number of beats per minute. Record the data in **Table 9**. Assign fitness points based on **Table 4** and record them in **Table 9**.

Table 2: Standing Pulse Rate

Pulse Rate (beats/min)	Fitness Points
< 71	3
71-80	3
81-90	2
91-100	1
101-110	1
111-120	0
121-130	0
≥ 131	-1

Test #3: Reclining Pulse Rate

Procedure:

1. The subject should recline for 5 minutes on a lab table/bench.
2. Determine the subject's resting pulse rate.
3. Count the number of beats for 30 seconds and multiply by 2. (NOTE: the subject should remain reclining for the next test.) The pulse rate is the number of beats per minute. Record the data in **Table 9**. Assign fitness points based on **Table 5** and record them in **Table 9**.

Table 5: Reclining Pulse Rate

Pulse Rate (beats/min)	Fitness Points
< 61	3
61-70	3
71-80	2
81-90	1
91-100	0
≥ 101	-1

Test #4: Baroreceptor Reflex (Pulse Rate Increase from Reclining to Standing)**Procedure:**

- The reclining subject should now stand up.
- Immediately* take the subject's pulse by counting the number of beats for 30 seconds. Multiply by 2 to determine the pulse rate in beats per minute. Record this value below. The observed increase in pulse rate is initiated by baroreceptors (pressure receptors) in the carotid artery and in the aortic arch. When the baroreceptors detect a drop in blood pressure they signal the medulla of the brain to increase the heartbeat and, consequently, the pulse rate.

Pulse immediately upon standing = _____ beats per minute

- Subtract the reclining pulse rate (recorded in Test 3) from the pulse rate immediately upon standing (recorded above for your subject) to determine the pulse rate increase upon standing. Record the data in **Table 9**. Assign fitness points based on **Table 6** and record them in **Table 9**.

Table 6: Pulse Rate Increase from Reclining to Standing

Reclining Pulse (beats/min)	Pulse Rate Increase on Standing (number of beats)				
	0-10	11-18	19-26	27-34	35-43
	Fitness Points				
< 61	3	3	2	1	0
61-70	3	2	1	0	-1
71-80	3	2	0	-1	-2
81-90	2	1	-1	-2	-3
91-100	1	0	-2	-3	-3
≥ 101	0	-1	-3	-3	-3

Test #5: Step Test – Endurance**Procedure:**

- The subject should do the following: Place your right foot on an 18-inch high stool. Raise your body so that your left foot comes to rest by your right foot. Return your left foot to the original position. Repeat this exercise 10 times, allowing 3 seconds for each step up.
- Immediately* after the completion of this exercise, measure the subject's pulse for 15 seconds and record below; measure again for 15 seconds (or 30 if the interval requires it) and record below; continue taking the subject's pulse and recording the rates at 60, 90, and 120 seconds.

Number of beats in the 0-15 second interval _____ X 4 = _____ beats per minute

Number of beats in the 16-30 second interval _____ X 4 = _____ beats per minute

Number of beats in the 31-60 second interval _____ X 2 = _____ beats per minute

Number of beats in the 61-90 second interval _____ X 2 = _____ beats per minute

Number of beats in the 91-120 second interval _____ X 2 = _____ beats per minute

- Observe the time that it takes for the subject's pulse rate to return to approximately the level that was recorded in Test 2. Record the data in **Table 9**. Assign fitness points based on **Table 7** and record them in **Table 9**.

Table 7: Time Required for Return of Pulse Rate to Standing Level after Exercise

Time (seconds)	Fitness Points
0-30	4
31-60	3
61-90	2
91-120	1
121+	1
1-10 beats above standing pulse rate	0
11-30 beats above standing pulse rate	-1

4. Subtract the subject's normal standing pulse rate (recorded in Test 2) from their pulse rate immediately after exercise (the 0-15 second interval) to obtain the pulse rate increase. Record the data in **Table 9**. Assign fitness points based on **Table 8** and record them in **Table 9**.

Table 8: Pulse Rate Increase after Exercise

Standing Pulse (beats/min)	Pulse Rate Increase Immediately After Exercise (beats)				
	0-10	11-20	21-30	31-40	41+
	Fitness Points				
< 71	3	3	2	1	0
71-80	3	2	1	0	-1
81-90	3	2	1	-1	-2
91-100	2	1	0	-2	-3
101-110	1	0	-1	-3	-3
111-120	1	-1	-2	-3	-3
121-130	0	-2	-3	-3	-3
≥ 131	0	-3	-3	-3	-3

Table 9: FITNESS DATA

	<u>Measurement</u>	<u>Points</u>
Test 1: Change in systolic pressure from reclining to standing	_____ mm Hg	
Test 2: Standing Pulse Rate	_____ beats/min	
Test 3: Reclining Pulse Rate	_____ beats/min	
Test 4: Baroreceptor Reflex—Pulse Rate Increase on Standing	_____ beats/min	
Test 5: Step Test Return on Pulse to Standing Rate after Exercise	_____ seconds	
Test 5: Step Test Pulse Rate Increase Immediately after Exercise	_____ beats/min	
	TOTAL SCORE	

Total Score	Relative Cardiac Fitness
18-17	Excellent
16-14	Good
13-8	Fair
7 or less	Poor

QUESTIONS and ANALYSIS:

1. Graph the temperature and heart rate data from the *Medaka* observations.


a. The independent variable is: _____

Use this to label the X-axis.

b. The dependent variable is: _____

Use this to label the Y-axis.

Title: _____



Name: _____

2. Why does temperature affect heart rate in ectothermic organisms?

3. Discuss what results you might obtain if you repeated this experiment two additional times using an endothermic organism starting with the **thermoneutral** zone of that species and then either increasing (one time) **or** decreasing (the other time) the environmental temperature.

4. Describe at least three different ways (metabolic or physical) that an **endothermic** organism can regulate its temperature.

5. Calculate Q_{10} for *Medaka* at **three** separate intervals during our investigation. SHOW YOUR WORK BELOW IN THE SPACES PROVIDED! SET UP THE CALCULATIONS!

For a 10° C interval, the heart rate Q_{10} can be calculated using sample data as follows:

$$Q_{10} = \frac{\text{rate at higher temperature}}{\text{rate at lower temperature}}$$

A general formula that can be used with **any** temperature interval is:

$$Q_{10} = \frac{k_2}{k_1}^{(10/t_2 - t_1)}$$

where:

t_2 = higher measurement temperature

t_1 = lower measurement temperature

k_2 = rate at temperature t_2

k_1 = rate at temperature t_1

Interval 1:

Interval 2:

Interval 3:

6. What do your calculations above tell you about the metabolic rates of *Medaka*?

Name: _____

7. Research and explain why blood pressure and heart rate differ when measured in a reclining position and in a standing position.

8. Research and explain why high blood pressure is a health concern.

9. Research and explain why an athlete must exercise harder or longer to achieve a maximum heart rate than a person who is not as physically fit.
